



# ***CP violation in Higgs decays***

**CFNS-SURGE 2025**

*A. I. Hernández-Juárez*

*alan.hernandezjua@alumno.buap.mx*

**June 6, 2025**

# The $H \rightarrow Z\gamma$ decay

PHYSICAL REVIEW LETTERS **132**, 021803 (2024)

Editors' Suggestion

Featured in Physics

## Evidence for the Higgs Boson Decay to a Z Boson and a Photon at the LHC

G. Aad *et al.*\*

(ATLAS and CMS Collaborations)



(Received 8 September 2023; accepted 27 November 2023; published 11 January 2024)

The first evidence for the Higgs boson decay to a Z boson and a photon is presented, with a statistical significance of 3.4 standard deviations. The result is derived from a combined analysis of the searches performed by the ATLAS and CMS Collaborations with proton-proton collision datasets collected at the CERN Large Hadron Collider (LHC) from 2015 to 2018. These correspond to integrated luminosities of around  $140 \text{ fb}^{-1}$  for each experiment, at a center-of-mass energy of 13 TeV. The measured signal yield is  $2.2 \pm 0.7$  times the standard model prediction, and agrees with the theoretical expectation within 1.9 standard deviations.

DOI: [10.1103/PhysRevLett.132.021803](https://doi.org/10.1103/PhysRevLett.132.021803)

# The $H \rightarrow Z\gamma$ decay

The signal strength is defined as

$$\mu_i^{Z\gamma} = \frac{\sigma_i \mathcal{B}^{Z\gamma}}{(\sigma_i)_{\text{SM}} (\mathcal{B}^{Z\gamma})_{\text{SM}}} = 2.2 \pm 0.7,$$

- $\sigma_i$  is the cross-section of the Higgs production
- $\mathcal{B}^{Z\gamma}$  is the branching ratio of the  $H \rightarrow Z\gamma$  decay

# The $H \rightarrow Z\gamma$ decay

The signal strength is defined as

$$\mu_i^{Z\gamma} = \frac{\sigma_i \mathcal{B}^{Z\gamma}}{(\sigma_i)_{\text{SM}} (\mathcal{B}^{Z\gamma})_{\text{SM}}} = 2.2 \pm 0.7,$$

Experimentally measured

SM prediction

- $\sigma_i$  is the cross-section of the Higgs production
- $\mathcal{B}^{Z\gamma}$  is the branching ratio of the  $H \rightarrow Z\gamma$  decay

# The $H \rightarrow Z\gamma$ decay

The signal strength is defined as

$$\mu_i^{Z\gamma} = \frac{\sigma_i \mathcal{B}^{Z\gamma}}{(\sigma_i)_{\text{SM}} (\mathcal{B}^{Z\gamma})_{\text{SM}}} = 2.2 \pm 0.7,$$

Possibilities of new physics!

- $\sigma_i$  is the cross-section of the Higgs production
- $\mathcal{B}^{Z\gamma}$  is the branching ratio of the  $H \rightarrow Z\gamma$  decay

# The $H \rightarrow Z\gamma$ decay

The signal strength is defined as

$$\mu_i^{Z\gamma} = \frac{\sigma_i \mathcal{B}^{Z\gamma}}{(\sigma)_{\text{SM}} (\mathcal{B}^{Z\gamma})_{\text{SM}}} = 2.2 \pm 0.7,$$


**Where is the new CP Violation?**

- $\mathcal{B}$  is the branching ratio of the  $H \rightarrow Z\gamma$  decay

new physics!

# The $H \rightarrow Z\gamma$ decay

The Higgs production is well measured and agrees with the SM production through a top quark loop in gluon fusion.



$$\sigma_i = (\sigma_i)_{\text{SM}}$$


Therefore, the new physics can only arise from the  $H \rightarrow Z\gamma$  decay:

$$\mu_i^{Z\gamma} = \frac{\mathcal{B}^{Z\gamma}}{\mathcal{B}_{\text{SM}}^{Z\gamma}} = 2.2 \pm 0.7,$$

A lot of models trying to explain this excess (THDM, MSSM, left-right models, new particles,...)

# The $H \rightarrow Z\gamma$ decay

The Higgs production is well measured and agrees with the SM production through a top quark loop in gluon fusion.



$$\sigma_i = (\sigma_i)_{\text{SM}}$$

Therefore, the new physics can only arise from the  $H \rightarrow Z\gamma$  decay:

$$\mu_i^{Z\gamma} = \frac{\mathcal{B}^{Z\gamma}}{\mathcal{B}_{\text{SM}}^{Z\gamma}} = 2.2 \pm 0.7,$$


CP violation can also explain this excess!

A lot of models trying to explain this excess (THDM, MSSM, left-right models, new particles,...)



# The $H \rightarrow Z\gamma$ decay

The  $H \rightarrow Z\gamma$  decay can be parametrized by the vertex function  $\Gamma_{Z\gamma H}^{\mu\nu}$ .



The general form of the vertex function  $\Gamma_{Z\gamma H}^{\mu\nu}$  is given as follows


$$\Gamma_{Z\gamma H}^{\mu\nu} = h_1^{Z\gamma} g^{\mu\nu} + \frac{1}{m_Z^2} \left\{ h_2^{Z\gamma} p_1^\nu p_2^\mu + h_3^{Z\gamma} \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{2\beta} \right\},$$

$$h_2^{Z\gamma} = \frac{2 m_Z^2}{m_Z^2 - m_H^2} h_1^{Z\gamma}.$$

Complex  
 $\sim 10^{-1}$  in the SM

# The $H \rightarrow Z\gamma$ decay

The  $H \rightarrow Z\gamma$  decay can be parametrized by the vertex function  $\Gamma_{Z\gamma H}^{\mu\nu}$ .



The general form of the vertex function  $\Gamma_{Z\gamma H}^{\mu\nu}$  is given as follows

$$\Gamma_{Z\gamma H}^{\mu\nu} = h_1^{Z\gamma} g^{\mu\nu} + \frac{1}{m_Z^2} \left\{ h_2^{Z\gamma} p_1^\nu p_2^\mu + h_3^{Z\gamma} \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{2\beta} \right\},$$

$$h_2^{Z\gamma} = \frac{2 m_Z^2}{m_Z^2 - m_H^2} h_1^{Z\gamma}.$$

Complex  
 $\sim 10^{-1}$  in the SM

CP-violating and zero  
 in the SM

# The $H \rightarrow Z\gamma$ decay

The signal strength  $\mu^{Z\gamma}$  can be expressed as follows

$$\mu^{Z\gamma} \simeq \frac{\mathcal{B}^{\mathbf{SM}}(H \rightarrow Z\gamma) + \delta\Gamma(H \rightarrow Z\gamma)/\Gamma_H}{\mathcal{B}^{\mathbf{SM}}(H \rightarrow Z\gamma)},$$


Where

$$\delta\Gamma(H \rightarrow Z\gamma) = g^2 \frac{(m_H^2 - m_Z^2)^3}{32 \pi m_H^3 m_Z^4} |h_3^{Z\gamma}|^2.$$

A. I. Hernández-Juárez, R. Gaitán and R. Martinez,  $H \rightarrow Z\gamma$  decay and CP violation, Phys. Rev. D 111, 015001 (2025), arXiv:2405.03094 [hep-ph].

$$\left| \text{Re}[h_3^{Z\gamma}] \right|, \left| \text{Im}[h_3^{Z\gamma}] \right| \lesssim 1.15 \text{ GeV} \quad \text{at 95 \% CL.}$$

First direct limit on  $h_3^{Z\gamma}$  from experimental data




# The $H \rightarrow Z\gamma$ decay

Effective Lagrangian that induces FCNC of the Higgs and  $Z$  boson:

$$\mathcal{L} = \frac{g}{c_W} \bar{f}_i \left( g_V^{ij} - g_A^{ij} \gamma^5 \right) f_j Z^\mu + \frac{g}{2m_W} \bar{f}_i \left( g_S^{ij} + g_P^{ij} \gamma^5 \right) f_j H,$$

$g_V^{ij}$ ,  $g_A^{ij}$ ,  $g_S^{ij}$  and  $g_P^{ij}$  complex constants

A possible new physics contribution:




It does not contribute to the  $H \rightarrow \gamma\gamma$  decay

# The $H \rightarrow Z\gamma$ decay

$$h_3^{Z\gamma} = \frac{g \, Q \, e \, m_Z^2 \, N_c}{4\pi^2 c_W m_W} \left\{ m_j \mathbf{C}_0 \left( 0, m_H^2, m_Z^2, m_j^2, m_j^2, m_i^2 \right) \left[ -\text{Im} \left\{ g_A^{ij} \left( g_S^{ij} \right)^* \right\} + \text{Im} \left\{ g_V^{ij} \left( g_P^{ij} \right)^* \right\} \right] \right. \\ \left. + m_i \mathbf{C}_0 \left( 0, m_H^2, m_Z^2, m_i^2, m_i^2, m_j^2 \right) \left[ \text{Im} \left\{ g_A^{ij} \left( g_S^{ij} \right)^* \right\} + \text{Im} \left\{ g_V^{ij} \left( g_P^{ij} \right)^* \right\} \right] \right\}$$

Calculated for the first time also

A possible new physics contribution:



It does not contribute to the  $H \rightarrow \gamma\gamma$  decay

# The $H \rightarrow Z\gamma$ decay

$$h_3^{Z\gamma} = \frac{g \, Q \, e \, m_Z^2 \, N_c}{4\pi^2 c_W m_W} \left\{ m_j \mathbf{C}_0 \left( 0, m_H^2, m_Z^2, m_j^2, m_j^2, m_i^2 \right) \left[ -\text{Im} \left\{ g_A^{ij} \left( g_S^{ij} \right)^* \right\} + \text{Im} \left\{ g_V^{ij} \left( g_P^{ij} \right)^* \right\} \right] \right. \\ \left. + m_i \mathbf{C}_0 \left( 0, m_H^2, m_Z^2, m_i^2, m_i^2, m_j^2 \right) \left[ \text{Im} \left\{ g_A^{ij} \left( g_S^{ij} \right)^* \right\} + \text{Im} \left\{ g_V^{ij} \left( g_P^{ij} \right)^* \right\} \right] \right\}$$

A. I. Hernández-Juárez, R. Gaitán and R. Martinez,  $H \rightarrow Z\gamma$  decay and CP violation, Phys. Rev. D 111, 015001 (2025), arXiv:2405.03094 [hep-ph].

**Limits on top quark FCNC couplings:**

$$|g_{V,A}^{tc}| < 0.0095, \quad |g_{S,P}^{tc}| \lesssim 0.25 \text{ GeV}$$

- We estimate that for FCNC of the top quark  $h_3^{Z\gamma} \approx 10^{-5}$ , too small to explain the  $\mu^{Z\gamma}$  excess.
- Contributions from new quarks are also possible and close to the bounds on  $h_3^{Z\gamma}$ .

# The $HZZ$ vertex

nature  
physics

ARTICLES


<https://doi.org/10.1038/s41567-022-01682-0>

Check for updates

OPEN

Measurement of the Higgs boson width and evidence of its off-shell contributions to  $ZZ$  production

The CMS Collaboration\*



$$m_H < 2m_Z$$



The Higgs boson must to be off-shell to produce two on-shell Z bosons



Submitted to: Phys. Lett. B.



CERN-EP-2023-03  
5th April 2023

Evidence of off-shell Higgs boson production from  $ZZ$  leptonic decay channels and constraints on its total width with the ATLAS detector


The ATLAS Collaboration

$$\Gamma_H = 3.2^{+2.4}_{-1.7} \text{ MeV}$$

The  $H \rightarrow ZZ^*$  well measured at the LHC

# The $HZZ$ vertex

Anomalous couplings for the  $ZZH$  vertex can be induced



$$= i \frac{g}{c_W} m_Z \Gamma_{\mu\nu}^{ZZH}(p_1^2, p_2^2, q^2)$$

Similar for the  $HW^\pm W^\mp$  case


$h_i^V$  in terms of the anomalous couplings

$$\Gamma_{\mu\nu}^{ZZH} = h_1^V g_{\mu\nu} + \frac{h_2^V}{m_Z^2} p_{1\nu} p_{2\mu} + \frac{h_3^V}{m_Z^2} \epsilon_{\mu\nu\alpha\beta} p_1^\alpha p_2^\beta,$$



# $HZZ$ vertex function

Anomalous couplings for the  $ZZH$  vertex can be also induced



Similar for the  $HW^\pm W^\mp$  case

$h_i^V$  in terms of the anomalous couplings


$$\Gamma_{\mu\nu}^{ZZH} = \underbrace{h_1^V}_{\text{CP-conserving}} \varepsilon_{\mu\nu} + \underbrace{\frac{h_2^V}{m_Z^2}}_{\text{CP-conserving}} p_{1\nu} p_{2\mu} + \underbrace{\frac{h_3^V}{m_Z^2}}_{\text{CP-violating}} \varepsilon_{\mu\nu\alpha\beta} p_1^\alpha p_2^\beta,$$

CP-conserving

CP-violating

# $HZZ$ vertex function

Anomalous couplings for the  $ZZH$  vertex can also be induced



Similar for the  $HW^\pm W^\mp$  case

$h_i^V$  in terms of the anomalous couplings


$$\Gamma_{\mu\nu}^{ZZH} = h_1^V g_{\mu\nu} + \frac{h_2^V}{m_Z^2} p_{1\nu} p_{2\mu} + \frac{h_3^V}{m_Z^2} \epsilon_{\mu\nu\alpha\beta} p_1^\alpha p_2^\beta,$$

Induced at one-loop level and complex in the SM.

¿Induced at three-loop level in the SM?  
 $h_3^V \sim 10^{-11}$

# $HZZ$ vertex function

Anomalous couplings for the  $ZZH$  vertex can also be induced



Similar for the  $HW^\pm W^\mp$  case

$h_i^V$  in terms of the anomalous couplings

$$\Gamma_{\mu\nu}^{ZZH} = h_1^V g_{\mu\nu} + \frac{h_2^V}{m_Z^2} p_{1\nu} p_{2\mu} + \frac{h_3^V}{m_Z^2} \epsilon_{\mu\nu\alpha\beta} p_1^\alpha p_2^\beta,$$

Induced at one-loop level and complex in the SM.

¿Induced at three-loop level in the SM?  
 $h_3^V \sim 10^{-11}$


Possibilities of new physics!

# $HZZ$ vertex function


Again FCNC of  $Z$  and  $H$  bosons: 
$$\mathcal{L} = \frac{g}{c_W} \bar{f}_i \left( g_V^{ij} - g_A^{ij} \gamma^5 \right) f_j Z^\mu + \frac{g}{2m_W} \bar{f}_i \left( g_S^{ij} + g_P^{ij} \gamma^5 \right) f_j H,$$

## Two different contributions

Type I:



Type II:




# $HZZ$ vertex function

Again FCNC of  $Z$  and  $H$  bosons: 
$$\mathcal{L} = \frac{g}{c_W} \bar{f}_i \left( g_V^{ij} - g_A^{ij} \gamma^5 \right) f_j Z^\mu + \frac{g}{2m_W} \bar{f}_i \left( g_S^{ij} + g_P^{ij} \gamma^5 \right) f_j H,$$

Two different contributions



Type I:



Calculated for the first time also

$h_3^V$  induced at the one-loop level

Type II:




# $HZZ$ vertex function

Again FCNC of  $Z$  and  $H$  bosons:  $\mathcal{L} = \frac{g}{c_W} \bar{f}_i \left( g_V^{ij} - g_A^{ij} \gamma^5 \right) f_j Z^\mu + \frac{g}{2m_W} \bar{f}_i \left( g_S^{ij} + g_P^{ij} \gamma^5 \right) f_j H,$

## Two different contributions



Type I:



$$h_3^V \sim 10^{-8}$$


Calculated for the first time also

Type II:



$$h_3^V \sim 10^{-7}$$

# $HZZ$ vertex function




- Small effects of  $h_3^H$  in the process  $gg \rightarrow H^* \rightarrow ZZ \rightarrow 4l$

¿Polarized process  $gg \rightarrow H^* \rightarrow Z_\lambda Z_\lambda \rightarrow 4l$  ?     $\lambda = R, L$  and  $0$

- Left-Right asymmetry:

$$\mathcal{A}_{LR}^H = \frac{\Gamma_{H^* \rightarrow Z_L Z_L} - \Gamma_{H^* \rightarrow Z_R Z_R}}{\Gamma_{H^* \rightarrow Z_L Z_L} + \Gamma_{H^* \rightarrow Z_R Z_R}}$$

# $HZZ$ vertex function



- Small effects of  $h_3^H$  in the process  $gg \rightarrow H^* \rightarrow ZZ \rightarrow 4l$


¿Polarized process  $gg \rightarrow H^* \rightarrow Z_\lambda Z_\lambda \rightarrow 4l$  ?  $\lambda = R, L$  and  $0$

- Left-Right asymmetry:

$$\mathcal{A}_{LR}^H \sim \text{Re}[h_1^H] \text{Im}[h_3^H] - \text{Re}[h_3^H] \text{Im}[h_1^H]$$



# $HZZ$ vertex function



- Small effects of  $h_3^H$  in the process  $gg \rightarrow H^* \rightarrow ZZ \rightarrow 4l$

¿Polarized process  $gg \rightarrow H^* \rightarrow Z_\lambda Z_\lambda \rightarrow 4l$ ?  $\lambda = R, L$  and  $0$

- Left-Right asymmetry:

$$\mathcal{A}_{LR}^H \sim \text{Re}[h_1^H] \text{Im}[h_3^H] - \text{Re}[h_3^H] \text{Im}[h_1^H]$$

$h_1^H$  complex in the SM

**CP-violation**

**Imaginary parts**

Similar asymmetries are possible in the  $Z^* \rightarrow Z_\lambda H$  and the  $HWW$  vertex

# Summary

- Effects of new physics are still possible in Higgs couplings.
- CP-violation can explain the reported excess in the  $H \rightarrow Z\gamma$  decay.
- New sources of CP-violation in the  $HZZ$  and  $HWW$ .

# ¡Gracias!



**¡Back up!**

# The $H \rightarrow Z\gamma$ decay

$$\Gamma(H \rightarrow Z\gamma) = g^2 \frac{m_H^2 - m_Z^2}{32 \pi m_H^3 m_Z^4} \left( 4 |h_1^{Z\gamma}|^2 m_Z^4 + |h_3^{Z\gamma}|^2 (m_H^2 - m_Z^2)^2 \right)$$

$$= \Gamma^{\mathbf{SM}}(H \rightarrow Z\gamma) + \delta\Gamma(H \rightarrow Z\gamma),$$

The general form of the vertex function  $\Gamma_{Z\gamma H}^{\mu\nu}$  is given as follows

$$\Gamma_{Z\gamma H}^{\mu\nu} = h_1^{Z\gamma} g^{\mu\nu} + \frac{1}{m_Z^2} \left\{ h_2^{Z\gamma} p_1^\nu p_2^\mu + h_3^{Z\gamma} \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{2\beta} \right\},$$



$$h_2^{Z\gamma} = \frac{2 m_Z^2}{m_Z^2 - m_H^2} h_1^{Z\gamma}.$$

Complex  
 $\sim 10^{-1}$

CP-violating and zero  
in the SM

# $HZZ$ vertex function

$$\mathcal{A}_{LR}^H \sim \text{Re}[h_1^H] \text{Im}[h_3^H] - \text{Re}[h_3^H] \text{Im}[h_1^H]$$



The observation of the  $\mathcal{A}_{LR}^H$  asymmetry would imply a new source of CP violation