

Unresolved Questions in Cold Nuclear Matter

C-J. Naim

Anaheim Convention Center, Anaheim (CA, USA)

**The 11th biennial workshop of the APS Topical Group on Hadronic Physics
(GHP2025)**

March 10, 2025



Proton-Proton Collisions

At large momentum transfer in pp, scale $Q \gg \Lambda_{\text{QCD}} \approx 200 \text{ MeV}$

$$pp \rightarrow \gamma^*/Z^0 \rightarrow \ell^+\ell^- + X \quad (\text{Drell-Yan})$$

Factorization of cross section = approximation

$$\frac{d\sigma_{pp}}{dy dQ} = \sum_{i,j} \int dx_1 f_i^p(x_1, \mu) \int dx_2 f_j^p(x_2, \mu) \frac{d\hat{\sigma}_{ij}(x_1, x_2, \mu')}{dy dQ} + \mathcal{O}\left(\frac{\Lambda_p^n}{Q^n}\right)$$

- ▶ $\hat{\sigma}_{ij}$: Partonic cross section calculable in perturbation theory
- ▶ x_1, x_2 : Fraction of momentum carried by the parton in the proton
- ▶ $f_{i,j}$: Parton Distribution Function (PDF), **universal**

Proton-Nucleus Collisions

Cross section in pA collisions assuming collinear factorization:

$$\frac{d\sigma_{pA}}{dy dQ} = \sum_{i,j} \int dx_1 f_i^p(x_1, \mu) \int dx_2 f_j^A(x_2, \mu) \frac{d\hat{\sigma}_{ij}(x_1, x_2, \mu')}{dy dQ} + \mathcal{O}\left(\frac{\Lambda_A^n}{Q^n}\right)$$

- ▶ Probing the **PDF of a nucleus** (without nuclear effects):

$$f_i^A = Z f_i^p + (A - Z) f_i^n$$


$$\sigma_{pA} = Z \sigma_{pp} + (A - Z) \sigma_{pn} \approx A \sigma_{pp}$$

- ▶ Investigating nuclear effects via:

$$R_{pA} \equiv \frac{1}{A} \frac{d\sigma_{pA}}{d\sigma_{pp}} \approx 1$$

Nuclear parton distribution functions (nPDF)

- ▶ EMC effect discovered in 1983 in DIS on nuclear targets
- ▶ **PDF is modified in nuclei** : $f_j^{p/A} \neq f_j^p$



- ▶ The nuclear modification factor depends on x

nPDF and data-sets

Historically, nPDFs were mainly extracted from DIS data


	EPS09	DSSZ	nCTEQ	EPPS16	EPPS21
e-DIS	✓	✓	✓	✓	✓
ν -DIS		✓		✓	✓
Drell-Yan pA	✓	✓	✓	✓	✓
RHIC hadrons	✓	✓	✓	✓	✓
LHC data pA (QED)				✓	✓
Drell-Yan π A				✓	✓
LHC data pA (D mesons)					✓

► Recent hA collision data included to:

- Extend the explored x range
- Access gluon nPDF more directly

→ **Possible biases from additional nuclear effects**

nPDF and data-sets




► nCTEQ15WZ and EPPS21 use heavy quark data in pA

→ Strongly impacting R_g^A

nPDF Scaling

$R_{pA}^{nPDF}(x, Q^2, \sqrt{s}) \equiv R_{pA}(x, Q^2)$ should scale as a function of \sqrt{s}



- Nuclear dependence for J/ψ and Drell-Yan production

Arleo Naïm Platchkov 1810.05120


- No scaling as a function of \sqrt{s} observed

→ **Exploring beyond nPDF effects!**

Exploring beyond nPDF effects

The nuclear medium affects hard processes differently.

- ▶ $hA \rightarrow \gamma^* + X$ (DY)
 - ▶ **Initial-state interactions**
- ▶ $eA \rightarrow e + h + X$ (SIDIS)
 - ▶ **Final-state interactions**




- ▶ $hA \rightarrow c\bar{c}(\rightarrow J/\psi) + X$ (Quarkonia)
 - ▶ **Initial- and final-state interactions**

How does the nuclear medium affect particle production?

Energy loss effects

Energy loss effects have successfully described nuclear data



- E866 and ALICE J/ψ suppression, $\langle \epsilon \rangle_{\text{FCEL}} \propto \sqrt{\hat{q}L}/M \cdot E$

Arleo Peigné [1204.4609](#), [1212.0434](#), Arleo Kolevaton Peigné Rustamova [2003.06337](#)


- E906 DY suppression, $\langle \epsilon \rangle_{\text{LPM}} \propto \alpha_s \hat{q} L^2$

Arleo Naïm Platchkov [1810.05120](#)

→ What about other effects?

Transverse Momentum Broadening


Broadening effects have successfully described nuclear data



- ▶ J/ψ , ψ' , Υ and DY data: a factor of 400 in beam energy!
- ▶ Broadening analysis reveals **universal scaling** across energies

nPDF including the energy loss effect

A global exhaustive fit: the (only) future path?



Avez Arleo [work ongoing](#)


- ▶ Global fit including nPDF and energy loss from J/ψ suppression
- ▶ Significant impact on the shadowing amplitude
 - Shadowing would be no more than 10-20% at $x \sim 10^{-5}$

Challenges in constraining gluon shadowing

Constraints on gluon shadowing from LHC pA data

- ▶ Limited experimental data for quarkonia and D-mesons
- ▶ Challenges in distinguishing shadowing from other nuclear effects
- ▶ Including energy loss in the global fit drastically reduces the shadowing amplitude by 10-20% at $x \sim 10^{-5}$


Ultra-peripheral collisions (UPC):



→ J/ψ production in UPCs to probe gluon shadowing

Shadowing amplitude

J/ψ production in UPCs: a direct probe of R_g



Guzey [CFNS cold QCD workshop \(2025\)](#)


- ▶ Strong nuclear suppression of coherent J/ψ photoproduction in Pb-Pb UPC@LHC due to large gluon shadowing at small x

Guzey Kryshen Strikman Zhalov [1305.1724](#), Guzey Zhalov [1307.4526](#)

- ▶ EPPS21, nCTEQ15HQ, nNNPDF3.0 use heavy quarks in pPb
→ A shadowing amplitude up to 60% at $x \sim 10^{-5}$

Nuclear absorption

Multiple scattering of $Q\bar{Q}$ bound state within the nucleons



- ▶ The typical size of a heavy nucleus is $L \sim 10$ fm
- ▶ J/ψ is mainly produced **outside** the nucleus at large y
 - No J/ψ absorption at LHC forward data

Energy loss or nuclear absorption?

The absorption of quarkonia remains an open question

- ▶ $\sigma_{\text{abs}}^{J/\psi} \sim 3 - 10 \text{ mb}$: extracted using pA at $y \sim 0$
→ **probably overestimated**

Laurenco Vogt Woehri [0901.3054](#), Arleo Tram [0612043](#)

- ▶ **Energy loss alone coherently explains J/ψ suppression in pA**
- ▶ Possible shadowing effects in nuclear matter: 10%, 20% or more?
- ▶ What remains of the role of absorption, $\sigma_{\text{abs}}^{J/\psi} \ll 3 \text{ mb}$?

→ Comparison: J/ψ suppression in eA vs pA collisions
The suppression should not be universal

Nuclear Data Challenges

Numerous nuclear data available, from fixed-target to LHC

- ▶ Difficult to interpret due to **multiple effects**
- ▶ Need to isolate specific effects through **golden observables**
- ▶ Importance of **global approaches** (global fits)
- ▶ Critical to estimate the **precise contribution of shadowing!**

→ The cold QCD effects are the primary source of uncertainties in the interpretation of AA collisions

Golden observables?

- ▶ $\mathcal{R} = R_{\text{pA}}^{J/\psi} / R_{\text{pA}}^{\psi} \sim S(\sigma_{\text{abs}}^{J/\psi}, L_A) / S(\sigma_{\text{abs}}^{\psi}, L_A)$
 - ▶ Mid rapidity region, small \sqrt{s}
 - ▶ Independent of shadowing: $Q_{J/\psi}^2 \sim Q_{\psi}^2$
 - ▶ Independent of FCEL: $\langle \epsilon \rangle_{\text{FCEL}} \propto 1/M_{\perp}$
- ▶ $\mathcal{R} = R_{\text{pA}}^{J/\psi} / R_{\text{pA}}^{\gamma}$
 - ▶ Weak shadowing dependence, strong sets correlations
 - ▶ Probe of the mass dependence of FCEL
- ▶ Transverse momentum broadening Δp_{\perp}^2 in eA and pA collisions
 - ▶ Independent of shadowing
 - ▶ Independent of energy loss
- ▶ J/ψ production
 - ▶ Test the non-universality of J/ψ suppression in eA and pA
 - ▶ Strong test of $\langle \epsilon \rangle_{\text{FCEL}}$ vs $\langle \epsilon \rangle_{\text{LPM}}$, + possible nuclear abs.
 - ▶ In eA, $\langle \epsilon \rangle_{\text{LPM}} \rightarrow 0$ at large \sqrt{s}

Key Questions in Nuclear Collisions

White paper in preparation:
Nuclear Cold QCD: Review and Future Strategy

1. **Energy Loss Mechanisms**

- ▶ Initial-state (DY), final-state (SIDIS)
- ▶ Initial/final-states (Quarkonia)

2. **Final-State Interactions**

- ▶ Nuclear absorption, comovers

3. **Shadowing vs. Saturation**

- ▶ Shadowing amplitude
- ▶ Distinction between leading-twist shadowing and gluon saturation

CFNS Cold QCD workshop

Global Insights and Future Directions

- ▶ **Nuclear data reveal a scaling violation** as a function of x
 - ▶ R_{pA} is not universal
 - ▶ Collinear factorization is not satisfied
- ▶ **Shadowing uncertainty** impacts all data interpretation
- ▶ **Energy loss** is key to describing the data
- ▶ **Strategy to address the three questions:**
 - ▶ Assess the limitations of hA data for nPDF studies
 - ▶ Enhance global fits by incorporating nuclear effects
 - ▶ Strongly constrain shadowing using future EIC DIS data
 - ▶ Identify and measure the key observables

No need for more data, but better data AND stronger collaboration between phenomenologists and experimentalists