

# First Gluon PDF from Large Momentum Effective Theory

William Good ([goodwil9@msu.edu](mailto:goodwil9@msu.edu))  
LaMET 2025  
10.08.2025

In Collaboration with Fei Yao, and Huey-Wen Lin  
Based on work: arXiv:2505.13321v2 (2025) (in review for PLB)



# Introduction and Background

Amaroso, *et al.* (Snowmass) *Acta Physica Polonica. B*, 53, 12 (2023)  
Aguilar, *et al.* *Eur. Phys. J. A* 55, 190 (2019).  
Achenbach, *et al.* (LRP) *Nucl. Phys. A* 1047:122874 (2024)



Image credit: CERN

# Introduction and Background

- Parton distribution functions (PDFs) provide the probability for a parton to carry a fraction  $x$  of the proton momentum, which is an important input for high energy scattering experiments

Amaroso, *et al.* (Snowmass) *Acta Physica Polonica. B*, 53, 12 (2023)  
Aguilar, *et al.* *Eur. Phys. J. A* 55, 190 (2019).  
Achenbach, *et al.* (LRP) *Nucl. Phys. A* 1047:122874 (2024)



Image credit: CERN

# Introduction and Background

- Parton distribution functions (PDFs) provide the probability for a parton to carry a fraction  $x$  of the proton momentum, which is an important input for high energy scattering experiments
- Phenomenological (pheno.) studies of the gluon PDF have some difficulties in obtaining the PDF in the large- $x$  region because of limited data

Amaroso, *et al.* (Snowmass) *Acta Physica Polonica. B*, 53, 12 (2023)  
Aguilar, *et al.* *Eur. Phys. J. A* 55, 190 (2019).  
Achenbach, *et al.* (LRP) *Nucl. Phys. A* 1047:122874 (2024)



Image credit: CERN

# Introduction and Background

- Parton distribution functions (PDFs) provide the probability for a parton to carry a fraction  $x$  of the proton momentum, which is an important input for high energy scattering experiments
- Phenomenological (pheno.) studies of the gluon PDF have some difficulties in obtaining the PDF in the large- $x$  region because of limited data
  - Ongoing and future experiments such as those at FNAL (HL-)LHC, LHeC, FCC(?), EIC, EicC will help reveal more information about the gluon PDFs

Amaroso, *et al.* (Snowmass) *Acta Physica Polonica. B*, 53, 12 (2023)  
Aguilar, *et al.* *Eur. Phys. J. A* 55, 190 (2019).  
Achenbach, *et al.* (LRP) *Nucl. Phys. A* 1047:122874 (2024)



Image credit: CERN

# Introduction and Background

- Parton distribution functions (PDFs) provide the probability for a parton to carry a fraction  $x$  of the proton momentum, which is an important input for high energy scattering experiments
- Phenomenological (pheno.) studies of the gluon PDF have some difficulties in obtaining the PDF in the large- $x$  region because of limited data
  - Ongoing and future experiments such as those at FNAL (HL-)LHC, LHeC, FCC(?), EIC, EicC will help reveal more information about the gluon PDFs
- The gluon PDF has been primarily studied through moments and the pseudo-PDF method, until last year, I presented tentative results for the gluon quasi-PDF from LaMET, without matching

Radyushkin PRD 96:034025 (2017)

Amaroso, *et al.* (Snowmass) Acta Physica Polonica. B, 53, 12 (2023)  
Aguilar, *et al.* Eur. Phys. J. A 55, 190 (2019).  
Achenbach, *et al.* (LRP) Nucl. Phys. A 1047:122874 (2024)



Image credit: CERN

# Introduction and Background

- Parton distribution functions (PDFs) provide the probability for a parton to carry a fraction  $x$  of the proton momentum, which is an important input for high energy scattering experiments
- Phenomenological (pheno.) studies of the gluon PDF have some difficulties in obtaining the PDF in the large- $x$  region because of limited data
  - Ongoing and future experiments such as those at FNAL (HL-)LHC, LHeC, FCC(?), EIC, EicC will help reveal more information about the gluon PDFs
- The gluon PDF has been primarily studied through moments and the pseudo-PDF method, until last year, I presented tentative results for the gluon quasi-PDF from LaMET, without matching
- With the new perturbative calculations for our operator, I can now present the results for the first gluon PDF from LaMET

Radyushkin PRD 96:034025 (2017)

Amaroso, *et al.* (Snowmass) *Acta Physica Polonica. B*, 53, 12 (2023)  
Aguilar, *et al.* *Eur. Phys. J. A* 55, 190 (2019).  
Achenbach, *et al.* (LRP) *Nucl. Phys. A* 1047:122874 (2024)




Image credit: CERN

# Gluon PDFs from LQCD


# Gluon PDFs from LQCD

- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams




# Gluon PDFs from LQCD

- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for LaMET extractions of gluon PDFs



# Gluon PDFs from LQCD

- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for LaMET extractions of gluon PDFs
- Most of the PDF extractions used the pseudo-PDF method:





# Gluon PDFs from LQCD

- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for

[Hadstruc] First nucleon gluon PDF from pseudo-PDF (2021)

of gluon PDFs

extractions used the pseudo-






# Gluon PDFs from LQCD

- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for

[Hadstruc] First nucleon gluon PDF from pseudo-PDF (2021)


[MSULat] First physical continuum nucleon gluon PDFs (2023)



# Gluon PDFs from LQCD


- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for

[Hadstruc] First nucleon gluon PDF from pseudo-PDF (2021)




PRD 104:094516(9) (2021)


[MSULat] First physical continuum nucleon gluon PDFs (2023)



PRD 108:014508(1), (2023)



[ETMC] Nucleon gluon PDF with singlet quark mixing (2023)




PRD 108:094515(9) (2023)

# Gluon PDFs from LQCD

- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for


[Hadstruc+JAM] Gluon helicity PDFs (2024)

PRD 109:036031(3), (2024)  
PRL 133:161901(16), (2024)




PRD 104:054516(9), (2021)


[JAM] First physical nucleon gluon PDFs (2023)



, (2023)




[ETMC] Nucleon gluon PDF with singlet quark mixing (2023)



PRD 108:094515(9), (2023)

# Gluon PDFs from LQCD




- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for



[Hadstruc+JAM] Gluon helicity PDFs (2024)  
 PRD 109:036031(3), (2024)  
 PRL 133:161901(16), (2024)


[MSULat] Updated kaon gluon PDF (2025)  
 arXiv:2506.03002 (in review for PRD)

neon gluon singlet quark



# Gluon PDFs from LQCD





- Gluon correlators are notoriously challenging in lattice QCD because they come only from noisy disconnected diagrams
- This limits the long distance signal important for



[Hadstruc+JAM] Gluon helicity PDFs (2024) PRD 109:036001 (2024) PRL 133:161901 (2024)


[MSULAT+JAM] Pion gluon data in a global fit arXiv:2507.22730 (in review for PRL)

...ated kaon ... nucleon gluon ... (25) ... nglet quark ... (2)



# Large Momentum Effective Theory


Ji, PRL 110:262002 (2013).  
Ji, Sci. China Phys. Mech. Astron. 57:1407 (2014).



# Large Momentum Effective Theory

Ji, PRL 110:262002 (2013).  
Ji, Sci. China Phys. Mech.  
Astron. 57:1407 (2014).

- The gluon PDF can be formally defined by the Fourier transform (FT) of gluon correlations along the light-cone




# Large Momentum Effective Theory

Ji, PRL 110:262002 (2013).  
Ji, Sci. China Phys. Mech.  
Astron. 57:1407 (2014).

- The gluon PDF can be formally defined by the Fourier transform (FT) of gluon correlations along the light-cone

$$g(x) = \int \frac{d\xi^-}{2\pi x P^+} e^{-ixP^+ \xi^-} \langle P | F_a^{+i}(\xi^-) \mathcal{W}(\xi^-, 0) F_a^{+i}(0) | P \rangle$$




# Large Momentum Effective Theory

Ji, PRL 110:262002 (2013).  
Ji, Sci. China Phys. Mech.  
Astron. 57:1407 (2014).

- The gluon PDF can be formally defined by the Fourier transform (FT) of gluon correlations along the light-cone

$$g(x) = \int \frac{d\xi^-}{2\pi x P^+} e^{-ixP^+\xi^-} \langle P | F_a^{+i}(\xi^-) \mathcal{W}(\xi^-, 0) F_a^{+i}(0) | P \rangle$$

- Instead, we measure Euclidean correlators in large momentum states, renormalize, Fourier transform, and match to the light-cone




# Large Momentum Effective Theory

Ji, PRL 110:262002 (2013).  
 Ji, Sci. China Phys. Mech. Astron. 57:1407 (2014).

- The gluon PDF can be formally defined by the Fourier transform (FT) of gluon correlations along the light-cone

$$g(x) = \int \frac{d\xi^-}{2\pi x P^+} e^{-ixP^+ \xi^-} \langle P | F_a^{+i}(\xi^-) \mathcal{W}(\xi^-, 0) F_a^{+i}(0) | P \rangle$$

- Instead, we measure Euclidean correlators in large momentum states, renormalize, Fourier transform, and match to the light-cone
- The gluon PDF has some freedom of operator choice. Until this work, these were the only two operators which had perturbative details for hybrid renormalization described explicitly in the literature:



$$O^{(1)}(z) = F^{zi}(z) W(z, 0) F_i^z(0)$$

$$O^{(2)}(z) = F^{z\mu}(z) W(z, 0) F_\mu^z(0)$$

Zhang, et al., PRL 121:142001 (2019)  
 Wang, et al. PRD 100:074509 (2019)  
 Yao, et al. JHEP 11(2023)021


# Large Momentum Effective Theory

Ji, PRL 110:262002 (2013).  
 Ji, Sci. China Phys. Mech. Astron. 57:1407 (2014).

- The gluon PDF can be formally defined by the Fourier transform (FT) of gluon correlations along the light-cone

$$g(x) = \int \frac{d\xi^-}{2\pi x P^+} e^{-ixP^+ \xi^-} \langle P | F_a^{+i}(\xi^-) \mathcal{W}(\xi^-, 0) F_a^{+i}(0) | P \rangle$$

- Instead, we measure Euclidean correlators in large momentum states, renormalize, Fourier transform, and match to the light-cone
- The gluon PDF has some freedom of operator choice. Until this work, these were the only two operators which had perturbative details for hybrid renormalization described explicitly in the literature:



$$O^{(1)}(z) = F^{zi}(z)W(z, 0)F_i^z(0)$$

$$O^{(2)}(z) = F^{z\mu}(z)W(z, 0)F_\mu^z(0)$$

Zhang, et al., PRL 121:142001 (2019)  
 Wang, et al. PRD 100:074509 (2019)  
 Yao, et al. JHEP 11(2023)021

- Previous pseudo-PDF studies have used this operator, which has better signal

$$O^{(3)}(z) = F^{ti}(z)W(z, 0)F_i^t(0) - F^{ij}(z)W(z, 0)F_{ij}(0)$$


Balitsky, et al., PLB 808:135621 (2020)

# (Almost) First GLuon PDF from LaMET

# (Almost) First Gluon PDF from LaMET

LaMET 2024


- Last year, we attempted the first gluon PDF from LaMET, but the first two operators provided very poor signal



# (Almost) First Gluon PDF from LaMET

LaMET 2024



- Last year, we attempted the first gluon PDF from LaMET, but the first two operators provided very poor signal
- On the other hand, the 3rd operator provided very good signal, but we had to make a guess on the renormalization:



# (Almost) First Gluon PDF from LaMET

LaMET 2024




- Last year, we attempted the first gluon PDF from LaMET, but the first two operators provided very poor signal
- On the other hand, the 3rd operator provided very good signal, but we had to make a guess on the renormalization:



# (Almost) First Gluon PDF from LaMET

LaMET 2024

- Last year, we attempted the first gluon PDF from LaMET, but the first two operators provided very poor signal
- On the other hand, the 3rd operator provided very good signal, but we had to make a guess on the renormalization:



# Some Help From an Expert

# Some Help From an Expert

- Without the perturbative QCD expertise in our group, we were dead in the water, until Dr. Fei Yao came to MSU for a seminar and we discussed this problem!

# Some Help From an Expert

- Without the perturbative QCD expertise in our group, we were dead in the water, until Dr. Fei Yao came to MSU for a seminar and we discussed this problem!
- We took the information from the pseudo-PDF matching paper for  $O^{(3)}(z)$  :  
to derive the Wilson coefficient and momentum space matching

Balitsky, *et al.*, PLB 808:135621 (2020)

# Some Help From an Expert

- Without the perturbative QCD expertise in our group, we were dead in the water, until Dr. Fei Yao came to MSU for a seminar and we discussed this problem!

Balitsky, *et al.*, PLB 808:135621 (2020)

- We took the information from the pseudo-PDF matching paper for  $O^{(3)}(z)$  : to derive the Wilson coefficient and momentum space matching

$$\mathcal{H}(z, \mu) = 1 + \frac{\alpha_s C_A}{2\pi} \left( \frac{5}{6} \ln \left( \frac{z^2 \mu^2 e^{2\gamma_E}}{4} \right) + \frac{3}{2} \right)$$

# Some Help From an Expert

- Without the perturbative QCD expertise in our group, we were dead in the water, until Dr. Fei Yao came to MSU for a seminar and we discussed this problem!

Balitsky, *et al.*, PLB 808:135621 (2020)


- We took the information from the pseudo-PDF matching paper for  $O^{(3)}(z)$  : to derive the Wilson coefficient and momentum space matching

$$\mathcal{H}(z, \mu) = 1 + \frac{\alpha_s C_A}{2\pi} \left( \frac{5}{6} \ln \left( \frac{z^2 \mu^2 e^{2\gamma_E}}{4} \right) + \frac{3}{2} \right)$$

- With these two pieces of information, we can properly renormalize our data for this operator and match to the light-cone!

# Lattice Information


Follana *et al.* PRD 75:054502, 2007.  
Bazavov, *et al.* [MILC], PRD 82:074501 2010.  
Bazavov, *et al.* [MILC], PRD 87:054505 2013.  
Bazavov, *et al.* [F Lattice and MILC] PRD 98:074512  
2018



# Lattice Information

Follana *et al.* PRD 75:054502, 2007.  
Bazavov, *et al.* [MILC], PRD 82:074501 2010.  
Bazavov, *et al.* [MILC], PRD 87:054505 2013.  
Bazavov, *et al.* [F Lattice and MILC] PRD 98:074512  
2018


- Used lattices with  $N_f = 2 + 1 + 1$  highly improved staggered quarks generated by the MILC collaboration with Wilson-clover fermions used in the valence sector



# Lattice Information

Follana *et al.* PRD 75:054502, 2007.  
Bazavov, *et al.* [MILC], PRD 82:074501 2010.  
Bazavov, *et al.* [MILC], PRD 87:054505 2013.  
Bazavov, *et al.* [F Lattice and MILC] PRD 98:074512  
2018


- Used lattices with  $N_f = 2 + 1 + 1$  highly improved staggered quarks generated by the MILC collaboration with Wilson-clover fermions used in the valence sector
- We use lattice spacing  $a \approx 0.12$  fm, and tune valence pion masses to  $M_\pi \approx 310$  and 690 MeV with volume  $24^3 \times 64$



# Lattice Information

Follana *et al.* PRD 75:054502, 2007.  
Bazavov, *et al.* [MILC], PRD 82:074501 2010.  
Bazavov, *et al.* [MILC], PRD 87:054505 2013.  
Bazavov, *et al.* [F Lattice and MILC] PRD 98:074512  
2018


- Used lattices with  $N_f = 2 + 1 + 1$  highly improved staggered quarks generated by the MILC collaboration with Wilson-clover fermions used in the valence sector
- We use lattice spacing  $a \approx 0.12$  fm, and tune valence pion masses to  $M_\pi \approx 310$  and 690 MeV with volume  $24^3 \times 64$
- $\sim 1.2$ M 2-point (2pt) correlator measurements over  $10^{13}$  configurations



# Lattice Information

Follana *et al.* PRD 75:054502, 2007.  
Bazavov, *et al.* [MILC], PRD 82:074501 2010.  
Bazavov, *et al.* [MILC], PRD 87:054505 2013.  
Bazavov, *et al.* [F Lattice and MILC] PRD 98:074512  
2018


- Used lattices with  $N_f = 2 + 1 + 1$  highly improved staggered quarks generated by the MILC collaboration with Wilson-clover fermions used in the valence sector
- We use lattice spacing  $a \approx 0.12$  fm, and tune valence pion masses to  $M_\pi \approx 310$  and 690 MeV with volume  $24^3 \times 64$
- $\sim 1.2$ M 2-point (2pt) correlator measurements over  $10^{13}$  configurations
- Gaussian momentum smearing to get signal up to 2.14 GeV boost momentum



# Lattice Information


Follana *et al.* PRD 75:054502, 2007.  
Bazavov, *et al.* [MILC], PRD 82:074501 2010.  
Bazavov, *et al.* [MILC], PRD 87:054505 2013.  
Bazavov, *et al.* [F Lattice and MILC] PRD 98:074512  
2018

- Used lattices with  $N_f = 2 + 1 + 1$  highly improved staggered quarks generated by the MILC collaboration with Wilson-clover fermions used in the valence sector
- We use lattice spacing  $a \approx 0.12$  fm, and tune valence pion masses to  $M_\pi \approx 310$  and 690 MeV with volume  $24^3 \times 64$
- $\sim 1.2$ M 2-point (2pt) correlator measurements over  $10^{13}$  configurations
- Gaussian momentum smearing to get signal up to 2.14 GeV boost momentum
- Two gauge link smearing sets for the 3-point (3pt) correlators: 5 steps of hypercubic smearing (HYP5) and Wilson flow with time  $t=3a^2$  (W3)



# Lattice Information

- Used lattices with  $N_f = 2 + 1 + 1$  highly improved staggered quarks generated by the MILC collaboration with Wilson-clover fermions used in the valence sector
- We use lattice spacing  $a \approx 0.12$  fm, and tune valence pion masses to  $M_\pi \approx 310$  and 690 MeV with volume  $24^3 \times 64$
- $\sim 1.2$ M 2-point (2pt) correlator measurements over  $10^{13}$  configurations
- Gaussian momentum smearing to get signal up to 2.14 GeV boost momentum
- Two gauge link smearing sets for the 3-point (3pt) correlators: 5 steps of hypercubic smearing (HYP5) and Wilson flow with time  $t=3a^2$  (W3)
- We look at the “strange” nucleon ( $N_s$ ), light nucleon ( $N_l$ ), strange pion ( $\eta_s$ ), and the light pion ( $\pi$ ) for renormalization and focus on the nucleons for the final results



# Hybrid-Ratio Scheme

Ji, *et al.* Nucl. Phys. B. 964:115311 (2021)

# Hybrid-Ratio Scheme

Ji, *et al.* Nucl. Phys. B. 964:115311 (2021)

- The hybrid-ratio scheme is a renormalization scheme which handles the linear divergence from the Wilson line self energy at long distances

# Hybrid-Ratio Scheme

Ji, *et al.* Nucl. Phys. B. 964:115311 (2021)

- The hybrid-ratio scheme is a renormalization scheme which handles the linear divergence from the Wilson line self energy at long distances
- We renormalize the quasi-PDF matrix elements as:

$$h^R(z, P_z) = \begin{cases} \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z,0)} & z \leq z_s \\ \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z_s,0)} \times e^{(\delta m + m_0)(z - z_s)} & z > z_s \end{cases}$$

# Hybrid-Ratio Scheme

Ji, *et al.* Nucl. Phys. B. 964:115311 (2021)

- The hybrid-ratio scheme is a renormalization scheme which handles the linear divergence from the Wilson line self energy at long distances
- We renormalize the quasi-PDF matrix elements as:

$$h^R(z, P_z) = \begin{cases} \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z,0)} & z \leq z_s \\ \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z_s,0)} \times e^{(\delta m + m_0)(z - z_s)} & z > z_s \end{cases}$$

- $z_s$  is a distance scale, before which the divergence is mostly ignorable

# Hybrid-Ratio Scheme

Ji, *et al.* Nucl. Phys. B. 964:115311 (2021)

- The hybrid-ratio scheme is a renormalization scheme which handles the linear divergence from the Wilson line self energy at long distances
- We renormalize the quasi-PDF matrix elements as:

$$h^R(z, P_z) = \begin{cases} \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z,0)} & z \leq z_s \\ \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z_s,0)} \times e^{(\delta m + m_0)(z - z_s)} & z > z_s \end{cases}$$

- $z_s$  is a distance scale, before which the divergence is mostly ignorable
  - Should not be much more than  $\sim 0.3$  fm

# Hybrid-Ratio Scheme

Ji, *et al.* Nucl. Phys. B. 964:115311 (2021)

- The hybrid-ratio scheme is a renormalization scheme which handles the linear divergence from the Wilson line self energy at long distances
- We renormalize the quasi-PDF matrix elements as:

$$h^R(z, P_z) = \begin{cases} \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z,0)} & z \leq z_s \\ \frac{h^B(0,0)}{h^B(0,P_z)} \frac{h^B(z,P_z)}{h^B(z_s,0)} \times e^{(\delta m + m_0)(z - z_s)} & z > z_s \end{cases}$$

- $z_s$  is a distance scale, before which the divergence is mostly ignorable
  - Should not be much more than  $\sim 0.3$  fm
- $\delta m + m_0$  can be fit by matching to the Wilson coefficients for the given operator:

$$(\delta m + m_0)z + I_0 \approx \ln \left[ \frac{\mathcal{H}(z, \mu)}{h^B(z, 0)} \right]$$


# $\delta m + m_0$ Fit Results

## $\delta m + m_0$ Fit Results

- Due to the coarse lattice spacing, we interpolate the data in the small- $z$  range. We do this, then fit to three points  $\{z-0.02 \text{ fm}, z, z +0.02 \text{ fm}\}$ , varying  $z$ . We used fixed  $\mu=2 \text{ GeV}$  for this first study


# $\delta m + m_0$ Fit Results

- Due to the coarse lattice spacing, we interpolate the data in the small- $z$  range. We do this, then fit to three points  $\{z-0.02 \text{ fm}, z, z+0.02 \text{ fm}\}$ , varying  $z$ . We used fixed  $\mu=2 \text{ GeV}$  for this first study



# $\delta m + m_0$ Fit Results


- Due to the coarse lattice spacing, we interpolate the data in the small- $z$  range. We do this, then fit to three points  $\{z-0.02 \text{ fm}, z, z +0.02 \text{ fm}\}$ , varying  $z$ . We used fixed  $\mu=2 \text{ GeV}$  for this first study



Different linear divergences for different smearing amounts is expected

# $\delta m + m_0$ Fit Results

- Due to the coarse lattice spacing, we interpolate the data in the small- $z$  range. We do this, then fit to three points  $\{z-0.02 \text{ fm}, z, z + 0.02 \text{ fm}\}$ , varying  $z$ . We used fixed  $\mu=2 \text{ GeV}$  for this first study




Statistical consistency across different hadrons, is a good sign!

Different linear divergences for different smearing amounts is expected

# $\delta m + m_0$ Fit Results

Due to the coarse lattice spacing, we interpolate the data in the small  $z$  range



0.30  
0.10 0.12

This is in strong contrast to the fits for the other operators in our previous study, which seem to have some hadron dependent contaminations

Statistical  
different

0.20 0.22 0.24

fits for different  
smearing amounts is expected

# Renormalized Nucleon Gluon Matrix Elements


# Renormalized Nucleon Gluon Matrix Elements

- Using the  $\delta m + m_0$ , we can renormalize the matrix elements (MEs), and also compare against “phenomenological matrix elements” obtained by matching the CT18 gluon PDF to the quasi-PDF and transforming to position space:

# Renormalized Nucleon Gluon Matrix Elements

- Using the  $\delta m + m_0$ , we can renormalize the matrix elements (MEs), and also compare against “phenomenological matrix elements” obtained by matching the CT18 gluon PDF to the quasi-PDF and transforming to position space:


WG, Yao, Lin, arXiv:2505.13321v2 (2025)



# Renormalized Nucleon Gluon Matrix Elements

- Using the  $\delta m + m_0$ , we can renormalize the matrix elements (MEs), and also compare against “phenomenological matrix elements” obtained by matching the CT18 gluon PDF to the quasi-PDF and transforming to position space:



WG, Yao, Lin, arXiv:2505.13321v2 (2025)



Reasonable consistency between the larger momentum lattice MEs and the pheno! Not concerned about low momentum MEs

# Light Nucleon MEs Are Similar, Just a Bit Noisier

WG, Yao, Lin, arXiv:2505.13321v2 (2025)



# Long Distance Extrapolation

- We picked the  $P_z = 1.71$  GeV matrix elements to fit the extrapolations because they had the best signal across the 4 sets of data

# Long Distance Extrapolation

- We picked the  $P_z = 1.71$  GeV matrix elements to fit the extrapolations because they had the best signal across the 4 sets of data
- We use a physically motivated extrapolation form to fit the data at long intermediate distances Gao, et al. Phys. Rev. Lett. 128, 142003 (2022)


$$h^R(z, P_z) \approx A \frac{e^{-m\nu}}{|\nu|^d}$$

# Long Distance Extrapolation

- We picked the  $P_z = 1.71$  GeV matrix elements to fit the extrapolations because they had the best signal across the 4 sets of data
- We use a physically motivated extrapolation form to fit the data at long intermediate distances Gao, et al. Phys. Rev. Lett. 128, 142003 (2022)

$$h^R(z, P_z) \approx A \frac{e^{-m\nu}}{|\nu|^d}$$

- Here, I show the extrapolation for W3 at the heavier pion mass as a best case




WG, Yao, Lin, arXiv:2505.13321v2 (2025)

# Long Distance Extrapolation

- We picked the  $P_z = 1.71$  GeV matrix elements to fit the extrapolations because they had the best signal across the 4 sets of data
- We use a physically motivated extrapolation form to fit the data at long intermediate distances Gao, et al. Phys. Rev. Lett. 128, 142003 (2022)


$$h^R(z, P_z) \approx A \frac{e^{-m\nu}}{|\nu|^d}$$

- Here, I show the extrapolation for W3 at the heavier pion mass as a best case
- The extrapolation fits the data fairly well and provides long distance matrix elements with much smaller error than the lattice




# Extrapolation on Other Data Sets

- The extrapolation works reasonably well on all 4 data sets,




# Extrapolation on Other Data Sets

- The extrapolation works reasonably well on all 4 data sets, though the strange nucleon HYP5 has converges more slowly than the rest



# Extrapolation on Other Data Sets


- The extrapolation works reasonably well on all 4 data sets, though the strange nucleon HYP5 has converges more slowly than the rest
- There may be some systematic effects from the fit ranges, which should be explored and controlled further



# Quasi-PDF and Light Cone Matching

# Quasi-PDF and Light Cone Matching


- We perform the Fourier transform to obtain the quasi-PDF



# Quasi-PDF and Light Cone Matching

- We perform the Fourier transform to obtain the quasi-PDF, then perform light-cone matching to get the PDF, ignoring mixing


$$x\tilde{g}(x, P^z, \mu) = \int_{-1}^1 dy F_{gg}(x, y, \mu) yg(y, \mu) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{(xP_z)^2}, \frac{\Lambda_{\text{QCD}}^2}{((1-x)P_z)^2}\right)$$



# Quasi-PDF and Light Cone Matching

- We perform the Fourier transform to obtain the quasi-PDF, then perform light-cone matching to get the PDF, ignoring mixing

$$x\tilde{g}(x, P^z, \mu) = \int_{-1}^1 dy F_{gg}(x, y, \mu) yg(y, \mu) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{(xP_z)^2}, \frac{\Lambda_{\text{QCD}}^2}{((1-x)P_z)^2}\right)$$




$\Lambda_{\text{QCD}} = 300 \text{ MeV}$ , as a conservative estimate

# Quasi-PDF and Light Cone Matching

- We perform the Fourier transform to obtain the quasi-PDF, then perform light-cone matching to get the PDF, ignoring mixing

$$x\tilde{g}(x, P^z, \mu) = \int_{-1}^1 dy F_{gg}(x, y, \mu) yg(y, \mu) + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{(xP_z)^2}, \frac{\Lambda_{\text{QCD}}^2}{((1-x)P_z)^2}\right)$$

- We find that the quasi-PDF has some negative region, but it is almost completely washed out by matching



$\Lambda_{\text{QCD}} = 300 \text{ MeV}$ , as a conservative estimate

# Light-Cone PDFs


Anderson, *et al.* arXiv 2501.00665 (2024)  
Hou, *et al.* (CTEQ) PRD 103(1):014013 (2021)  
Cerutti, *et al.* PRD 111:094013 (2025)  
WG, Yao, Lin, arXiv:2505.13321v2 (2025)

# Light-Cone PDFs

Anderson, *et al.* arXiv 2501.00665 (2024)  
Hou, *et al.* (CTEQ) PRD 103(1):014013 (2021)  
Cerutti, *et al.* PRD 111:094013 (2025)  
WG, Yao, Lin, arXiv:2505.13321v2 (2025)

- We compare the PDFs across the 4 data sets,

WG, Yao, Lin, arXiv:2505.13321v2 (2025)




# Light-Cone PDFs

Anderson, *et al.* arXiv 2501.00665 (2024)  
Hou, *et al.* (CTEQ) PRD 103(1):014013 (2021)  
Cerutti, *et al.* PRD 111:094013 (2025)  
WG, Yao, Lin, arXiv:2505.13321v2 (2025)

- We compare the PDFs across the 4 data sets,

WG, Yao, Lin, arXiv:2505.13321v2 (2025)





The W3 PDFs are more negative, suggesting that the smearing is affecting the physics at this level

# Light-Cone PDFs

Anderson, *et al.* arXiv 2501.00665 (2024)  
Hou, *et al.* (CTEQ) PRD 103(1):014013 (2021)  
Cerutti, *et al.* PRD 111:094013 (2025)  
WG, Yao, Lin, arXiv:2505.13321v2 (2025)

- We compare the PDFs across the 4 data sets, and compare the two HYP5 results against phenomenological PDFs

WG, Yao, Lin, arXiv:2505.13321v2 (2025)





The W3 PDFs are more negative, suggesting that the smearing is affecting the physics at this level

# Light-Cone PDFs

Anderson, *et al.* arXiv 2501.00665 (2024)  
Hou, *et al.* (CTEQ) PRD 103(1):014013 (2021)  
Cerutti, *et al.* PRD 111:094013 (2025)  
WG, Yao, Lin, arXiv:2505.13321v2 (2025)

- We compare the PDFs across the 4 data sets, and compare the two HYP5 results against phenomenological PDFs

WG, Yao, Lin, arXiv:2505.13321v2 (2025)



The W3 PDFs are more negative, suggesting that the smearing is affecting the physics at this level

The HYP5 PDFs are most consistent with the CT18 PDF, and only a standard deviation or two from the other PDFs

# Outlook and Conclusion

# Outlook and Conclusion

- We have obtained the first gluon PDFs from LaMET at a heavier than physical pion mass and a somewhat coarse lattice spacing

# Outlook and Conclusion

- We have obtained the first gluon PDFs from LaMET at a heavier than physical pion mass and a somewhat coarse lattice spacing
- We find currently that smearing has some effect on the physics, which will need to be explored in more detail

# Outlook and Conclusion

- We have obtained the first gluon PDFs from LaMET at a heavier than physical pion mass and a somewhat coarse lattice spacing
- We find currently that smearing has some effect on the physics, which will need to be explored in more detail
- We still need to understand lattice spacing effects, along with eventual handling of perturbative uncertainties with leading renormalon resummation and renormalization group resummation

# Outlook and Conclusion

- We have obtained the first gluon PDFs from LaMET at a heavier than physical pion mass and a somewhat coarse lattice spacing
- We find currently that smearing has some effect on the physics, which will need to be explored in more detail
- We still need to understand lattice spacing effects, along with eventual handling of perturbative uncertainties with leading renormalon resummation and renormalization group resummation
- Reducing statistical uncertainties is still a major challenge, as well


# Outlook and Conclusion

- We have obtained the first gluon PDFs from LaMET at a heavier than physical pion mass and a somewhat coarse lattice spacing
- We find currently that smearing has some effect on the physics, which will need to be explored in more detail
- We still need to understand lattice spacing effects, along with eventual handling of perturbative uncertainties with leading renormalon resummation and renormalization group resummation
- Reducing statistical uncertainties is still a major challenge, as well
- Alex will show her results for a similar study with various lattice spacings with better control of discretization effects using self-renormalization

Backup Slides

# $\delta m + m_0$ Interpolation Order Dependence


- We checked how the interpolation affects the  $\delta m + m_0$  fit, and we see convergence by order cubic interpolation



$N_s$  W3

# Extrapolation Fit Range Dependence (1)

- The errors on the extrapolation are quite sensitive to the fit range, while the central values are statistically indistinguishable
- Here we pick a very short range vs a much longer range for the extrapolations



$N_s$  W3

# Extrapolation Fit Range Dependence

- These errors do have noticeable effects on the final PDF, as well
- We think this could be a significant source of systematic uncertainty at the current level of statistics, so in the future, we would like to consider model averaging or some more sophisticated method to do the Extrapolations
- Currently, we just use an intermediate fit range

